

RAUNDS URBAN
DISTRICT COUNCIL

ANNUAL REPORT
of the
MEDICAL OFFICER OF HEALTH
A. MCINNES, M.B., D.P.H.

1 9 6 3



R A U N D S U R B A N
D I S T R I C T C O U N C I L

A N N U A L R E P O R T
of the
MEDICAL OFFICER OF HEALTH
A. MCINNES, M.B., D.P.H.

1 9 6 3

Chairman, 1963: Mrs. R. Gardner
Clerk: B.M. Killick

Medical Officer of Health: A. McInnes, M.B., D.P.H.

Public Health Inspector
and Surveyor: G. Whittam, F.I.A.S., M.R.S.I.

Population: 4,620

		Dwellings	Population	Rateable Value £	Penny Rate £. s. d		
1958	...	1,621	4,650	35,572	136.	6.	0.2
1959	...	1,615	4,680	39,693	139.	12.	3.9
1960	...	1,637	4,670	40,305	157.	8.	3.77
1961	...	1,650	4,570	40,886	161.	1.	6.42
1962	...	1,669	4,610	41,524	162.	8.	8.99
1963	...	1,681	4,620	116,629	164.	15.	5.16

Census Returns

			Population	No. of Houses	No. per house
1901	...	...	3,811)	847	4.5
1911	...	...	3,874	850	4.5
1921	...	...	3,761	920	4.08
1931	...	...	3,687)	1,012	3.65
1941	...	...	(5,392 Raunds	1,395	3.86
1951	...	...	(4,579 and	1,512	3.02
1961	...	...	(4,570 Stanwick	1,650	2.77

			Rounds	Stanwick
1801	...	...	800	332
1831	...	...	1,370	503
1841	...	...	1,653	577
1851	...	...	1,873	609
1861	...	...	2,337	669
1877	...	...	2,580	678

[illegible]

Mr. Chairman,

In this report for 1963 some of the tables given in the report for 1962 are repeated. The statistics, over a period of years, give a sense of perspective and may indicate a trend. But in dealing with a small numerical basis, such as the population of Raunds, statistics must not lead to the general conclusion from the part to the whole, from Raunds to the whole of England and Wales. Nevertheless, statistics show that Raunds is a healthy district with a more than average aged population. Indeed the figures might suggest that Raunds is one third Industrial, one third Agricultural and one third a Health Resort. It should be noted that in 1963 deaths exceed births by 18% and that the crude death rate exceeds that of the County by almost 28% and that of England and Wales by 36%. This relatively high death rate is not due to disease, but to age. The table at the end of this report shows that death over 75 years were 54%, those over 80 years 34.7% and those over 90 years 8½% of the total deaths in 1963. And to the same point the birth rate in Raunds in 1963 was only 72% of the County and 70% of that of England and Wales.

Birth Rate

The number of births and a series of rates are given below. Up to 1950 only crude Birth Rates could be given, but for 1950 and afterwards a comparability factor has been issued so that standard Birth Rate = crude Birth Rate x comparability factor. For Raunds the comparability factor for 1959, 1960, 1961, 1962 was 1.16 and in 1963 was 1.2. This relatively high comparability factor suggests a population above the average in age.

Live Births

TOTAL LIVE BIRTHS in Raunds Urban District:-

				1960		1961		1962		1963	
				M	F	M	F	M	F	M	F
Legitimate	...	...	...	21	35	35	30	33	27	28	32
Illegitimate		...	...	0	0	0	2	0	1	1	0
TOTAL	...	...	...	21	35	35	32	33	28	29	32

Illegitimate rate per 1,000 Live Births 0.0 29.87 16.4 16.4

BIRTH RATES per 1,000 of population:-

Raunds U.D.C. - Crude	...	12.0	14.66	13.2	13.2
Standard	...	13.92	17.4	15.3	16.0
England and Wales - Crude	...	17.1	17.4	18.0	18.2
Administrative County - Crude		17.7	18.04		18.7

Still Births

A still birth is defined as the issue of a dead child after twenty-eight weeks of pregnancy. The two still births were in hospital.

				1960		1961		1962		1963	
				M	F	M	F	M	F	M	F
Legitimate	...	...	...	0	0	1	0	0	1	0	2
Illegitimate		...	...	0	0	0	0	0	0	0	0
TOTAL	...	...	...	0	0	1	0	0	1	0	2

Digitized by the Internet Archive
in 2018 with funding from
Wellcome Library

Rate per 1,000 of Live and Still Births:-				1960	1961	1962	1963
Raunds Urban District	...	...		0.0	15.0	16.1	31.7
England and Wales	...	...		19.7	19.1	18.1	17.2
Administrative County	...	...		16.32	17.6		14.85

Deaths of Children under 1 year

				1960		1961		1962		1963	
				M	F	M	F	M	F	M	F
Legitimate	...	...	...	1	0	0	3	1	0	0	1
Illegitimate	...	...	...	0	0	0	0	0	0	0	0
TOTAL	...	...	...	1	0	0	3	1	0	0	1

Rate per 1,000 of Live Births				1960	1961	1962	1963
Raunds Urban District	...	...		17.86	46.0	16.1	16.4
England and Wales	...	...		21.7	21.6	21.4	21.1
Administrative County	...	...		22.57	17.61		17.7

Deaths of Children under 4 weeks

				1961		1962		1963	
				M	F	M	F	M	F
Legitimate	...	...	...	0	2	1	0	0	1
Illegitimate	...	...	...	0	0	0	0	0	0
TOTAL	...	...	...	0	2	1	0	0	1

Rate per 1,000 of Live Births				1963
Raunds Urban District	...	...		16.4
England and Wales	...	...		14.2

Deaths of Children under 1 week

				1962		1963	
				M	F	M	F
Legitimate	...	...	...	1	0	0	1
Illegitimate	...	...	...	0	0	0	0
				1	0	0	1

Rate per 1,000 of Live Births				1963
Raunds Urban District	...	...		16.4
England and Wales	...	...		12.1

Peri-Matal Mortality (Still Births and Deaths under 1 Week)

Rate per 1,000 of Total Live and Still Births				1963
Raunds Urban District	...	...		48.0
England and Wales	...	...		29.3

This division of Infantile Mortality into under a week, under a month and under a year is an attempt to separate statistically inherent causes of death from social causes. This child died in hospital very soon after birth.

Maternal Mortality

	1960	1961	1962	1963
	0	0	0	0
Rate per 1,000 Live and Still Births:-				
Raunds Urban District	0.0	0.0	0.0	0.0
England and Wales	0.39	0.33		0.28
Administrative County	0.37	0.55		0.20

There has been no maternal death since 1935, the date of the inclusion of Stanwick in the district.

Of the 243 mothers who died in England and Wales in 1963, 60 died of sepsis and the other 183 died from complications of pregnancy or childbirth in the lying in period. In 1963 the chances of death in childbirth were 1:2500 births, whereas in 1926 the chances were 1:226. In the Scandinavian countries and in the U.S.A. the chances are similar to those in England and Wales. In Italy and Portugal the chances are 1:870; in Japan 1:760 and in Ceylon 1:330. The great reduction in maternal mortality is due first to the sulphonamide drugs and later to Pencillin and other anti-biotics in the treatment of and the prevention of sepsis (usually post-natal). The reduction in the other causes of maternal mortality is due to the meticulous ante-natal care. Before the twenties and thirties of this century, medical interest in the pregnant mother was very little and nature was allowed to take its course for good or evil.

Death Rate

Below are given the number of deaths and a table of death rates per 1,000 of population. A Comparability Factor has been given so that Crude Death Rate x Comparability Factor = Standard Death Rate. The necessity of this factor for the purposes of comparison is due to an unequal distribution of age groups and also, to a lesser degree, of the sexes. For example: Bournemouth and Cheltenham are more likely to have a greater number in the older age groups than say Coventry or Wigan, where most are of the earning ages. Females have a greater expectation of life than males. This also allows for the presence in a district of institutions especially for the elderly. You may have noticed that deaths exceed births, a natural decrease of 72 - 61 = 11, or .24% of total population.

A classification of the causes of death with sex and age is given in tables at the end of the report.

Number of Deaths

	1960	1961	1962	1963
Males	35	23	27	34
Females	<u>21</u>	<u>35</u>	<u>37</u>	<u>38</u>
	<u>56</u>	<u>58</u>	<u>64</u>	<u>72</u>

DEATH RATE:-

		1960	1961	1962	1963
Raunds Urban District -	Crude ...	12.0	12.7	13.88	15.6
	Standard	10.3	10.53	11.8	13.1
England and Wales ...		11.5	12.0	11.9	12.2
Administrative County		10.88	11.18		11.2
Comparability Factor 1958 = 0.85)					
"	"	1959 = 0.84)	These comparability factors indicate a more aged population than the average of England and Wales.		
"	"	1960 = 0.86)			
"	"	1961 = 0.83)			
"	"	1962 = 0.85)			
"	"	1963 = 0.84)			

The total Death returns of a public health district are made up of (1) Residents who die in the district and (2) Residents who die outside the district and their death transferred inwardly to the home district. Below are two tables for periods of 10 and 11 years each, (1) After the National Health Service Act and (2) Before the inception of the Act. Although the figures involved are small there is consistent evidence that people are living longer and also that the number of inwardly transferable deaths is increasing in proportion to the total deaths. Most of these inward transfers are of elderly people dying in institutions outside the district. This increase in the proportion of transferable deaths suggests that the family as the basis of social life is giving way to that of the community. The old people are without doubt well treated in the State institutions but nevertheless there is a severance of a link.

Year	After N.H.S.				Year	Before N.H.S.			
	Total Deaths	No. over 70	Rate	Inward Transfers Rate		Total Deaths	No. over 65	Rate	Inward Transfers Rate
1953	41	26	63%	16 39%	1916	48	18	37%	7 16%
1954	44	30	70%	19 43%	1917	41	18	44%	4 10%
1955	81	45	60%	23 28%	1918	54	17	31%	7 13%
1956	56	39	70%	24 43%	1919	40	26	50%	5 12.5%
1957	52	33	63%	12 23%	1920	45	22	50%	8 18%
1958	52	34	65%	20 39%	1921	39	21	54%	4 10%
1959	62	34	55%	17 28%	1922	34	12	35%	0 0%
1960	56	30	54%	18 32%	1923	46	20	44%	4 9%
1961	58	39	67%	23 40%	1924	41	18	44%	7 17%
1962	64	43	67%	32 50%	1925	32	14	44%	11 30%
1963	72	44	61%	33 45.8%					

It is to be noted that from 1953 the number is over 70 years and from 1916 the number is over 65 years. The expectation of life is distinctly better from 1953 and this table also shows distinctly more inward transfers.

Infectious Diseases

Scarlet Fever

14 cases were notified of a very mild type. This disease is now generally so mild that a diagnosis can be difficult.

Erysipelas

None.

Pneumonia

None

Typhoid and Paratyphoid

None

Cerebro-Spinal Fever

None

Measles

There were 45 notifications.

Whooping Cough

There were 3 notifications.

Acute Poliomyelitis and Polio-encephalitis

None

Enteric Fever

None

Food Poisoning

None

Influenza

None

Puerperal Pyrexia

None

Tuberculosis

One case of Pulmonary tuberculosis was notified in 1963.

Tuberculosis - Number on Register

The number of cases of Tuberculosis on the Register during the past eleven years was as follows:-

			<u>Respiratory</u>	<u>Non-respiratory</u>
31st December, 1953	...	...	24	5
31st December, 1954	...	...	29	7
31st December, 1955	...	...	22	5
31st December, 1956	...	...	22	5
31st December, 1957	...	...	24	6
31st December, 1958	...	...	18	4
31st December, 1959	...	...	15	4
31st December, 1960	...	...	11	5
31st December, 1961	...	...	10	4
31st December, 1962	...	...	9	4
31st December, 1963	...	...	5	4

Toxic Chemicals in Agriculture

Toxic Chemicals in Agriculture

Di-Nitro compounds. The chief are DNC and DNBP and are mostly used for weed killing from the end of March until the middle of June and most commonly for corn crops and for the destruction of potato haulm, usually in August and September. Farm workers are likely to suffer.

DNC is a cumulative poison for which there is no known antidote. When the concentration in the blood of the worker is well below 20 micro-grammes per gramme of blood the effect is to produce an exaggerated feeling of well-being. When the concentration approaches the 20 mark, the feeling of well-being is replaced by a feeling of fatigue, thirst and sweating. When a concentration of over 20 is reached the patient may be seriously ill.

Workers with DNC and DNBP should be under medical supervision for blood testing.

In this district blood may be tested at

1. Department of Pathology,
Memorial Hospital,
Peterborough.
2. Department of Biochemistry,
Addenbrooks Hospital,
Cambridge.
3. Department of Pathology,
Royal Infirmary,
Leicester.

Water Supply

Raunds is supplied with water by the Nene and Ouse Water Board and the following is an analysis of a sample of water from the Raunds Meadows supply:-

Sample of water labelled "Raw Water, Raunds Well (Pumping Station at Meadow Lane)" received on the 9th October, 1963 from Mr. H.B. Wilson, Nene & Ouse Water Board.

Physical Examination

Clear and bright, odourless and colourless (Hazen - less than 5.0)

General Chemical Examination

Reaction 7.0

					<u>Parts per 100,000</u>
Free Carbon Dioxide (CO_2)	...	...	...	...	2.01
Ammoniacal Nitrogen (N) ²	...	...	...	...	0.0029
Albuminoid Nitrogen (N)	...	...	...	...	0.0085
Nitrous Nitrogen	...	...	...	...	absent
Nitric Nitrogen (N)	...	...	...	...	0.40
Hardness CaCO_3 Clark)	...	...	...	...	40.0
Temporary	...	...	...	...	21.0
Permanent	...	...	...	...	19.0
Oxygen absorbed in 3 hours at 37°C	...	...	...	...	0.0660
Alkalinity (CaCO_3)	...	...	...	...	29.25
Total Solids	...	...	...	...	86.9
Poisonous Metals					absent

<u>Mineral Analysis</u>					<u>Parts per 100,000</u>
Calcium (Ca)	...	...	...	...	16.25
Magnesium (Mg)	...	...	...	...	1.15
Carbonate (CO ₃)	...	...	...	...	17.6
Chloride (Cl)	...	...	...	...	8.7
Sulphate (SO ₄)	...	...	...	...	15.13
Nitrate (NO ₃)	...	...	...	...	1.77
Iron	...	...	...	...	absent
Sodium	...	...	...	...	11.0

Microscopical Examination of Deposit ... None

Plumbo-Solvency - No action on lead in 3 days at 21°C

Bacteriological Examination

Coliform organisms absent in 100 mls.

Number of microorganisms per ml developing at 37°C = nil

Number of microorganisms per ml developing at 21°C = 12

R E M A R K S

The results obtained on the analysis of this sample indicate a hard water containing little organic matter and relatively few bacteria.

I am of opinion that this water as evidenced by the sample is fit for drinking purposes.

S. GREENBURGH

24th October, 1963.

Public Analyst.

Stanwick receives most of its water supply from Woodford source. An analysis of a sample from this supply is given herewith:-

Sample of water labelled "Raw Water, Well at Woodford Pumping Station" received on the 9th October, 1963 from Mr. H.B. Wilson, Nene & Ouse Water Board.

Physical Examination

Very slight deposit, odourless and colourless (Hazen - less than 5.0)

General Chemical Examination

Reaction pH 7.2

					<u>Parts per 100,000</u>
Free Carbon Dioxide (CO ₂)	...	...	...	...	1.58
Ammoniacal Nitrogen (N) ²	...	...	...	...	nil
Albuminoid Nitrogen (N)	...	...	...	...	0.0036
Nitrous Nitrogen	...	...	...	...	absent
Nitric Nitrogen (N)	...	...	...	...	0.35
Hardness (CaCO ₃ Clark)	...	...	...	...	35.8
Temporary	...	...	...	...	17.1
Permanent	...	...	...	...	18.7
Oxygen absorbed in 3 hours at 37°C	...	...	...	...	0.0412
Alkalinity (CaCO ₃)	...	...	...	...	24.75
Total Solids	...	...	...	...	75.9
Poisonous Metals	...	...	...	...	absent

Mineral AnalysisParts per 100,000

Calcium (Ca)	...	...	...	...	14.86
Magnesium (Mg)	...	...	...	...	0.82
Carbonate (CO ₃)	...	...	...	...	14.95
Chloride (Cl)	...	...	...	...	6.9
Sulphate (SO ₄)	...	...	...	...	15.46
Nitrate (NO ₃)	...	...	...	...	1.55
Iron	...	...	...	...	trace
Sodium	...	...	...	...	13.76

Microscopical Examination of Deposit

Mainly mineral matter.

Plumbo-Solvency - No action on lead in 3 days at 21°C

Bacteriological Examination

Coliform organisms absent in 100 mls.

Number of microorganisms per ml developing at 37°C = nil

Number of microorganisms per ml developing at 21°C = 5

R E M A R K S

The results obtained on the analysis of this sample indicate a hard water containing little organic matter and few bacteria.

I am of opinion that this water as evidenced by the sample is fit for drinking purposes.

S. GREENBURGH

24th October, 1963.

Public Analyst.

R E P O R T
on the

Bacteriological Examination of the undermentioned samples of Water received on the 18th August 1964 from Mr. H.B. Wilson, Engineer, Nene and Ouse Water Board, Thrapston House, Thrapston, Northants.

<u>SAMPLE</u>	B.Coli in 100 mls	Number of microorganisms per mil at 37°C	Number of microorganisms per mil at 21°C
RAUNDS	absent	nil	nil
STANWICK	absent	nil	1

A. McINNES

Medical Officer of Health.

STATISTICAL TABLES

CAUSES OF DEATH

There were no deaths in the years 1 - 5, 5 - 15, 15 - 25, 25 - 35.

Causes of Death		Sex	Total All Ages	Under 4 weeks	4 weeks and under 1 year	35 -	45 -	55 -	65 -	75 and over
10. Malignant Neoplasm, Stomach	M	2	-	-	-	-	-	1	-	1
	F	-	-	-	-	-	-	-	-	-
11. Malignant Neoplasm, Lung, Bronchus	M	1	-	-	-	-	-	-	-	1
	F	-	-	-	-	-	-	-	-	-
12. Malignant Neoplasm, Breast	M	-	-	-	-	-	-	-	-	-
	F	1	-	-	-	-	1	-	-	-
14. Other Malignant and Lymphatic Hooplasms	M	3	-	-	-	-	-	1	1	1
	F	3	-	-	-	-	1	2	-	-
16. Diabetes	M	1	-	-	-	-	-	-	-	1
	F	-	-	-	-	-	-	-	-	-
17. Vascular Lesions of Nervous System	M	2	-	-	-	-	-	1	-	1
	F	9	-	-	-	-	-	2	2	5
18. Coronary Disease, Angina	M	6	-	-	-	-	-	1	1	4
	F	7	-	-	-	-	-	-	2	5
19. Hypertension with Heart Disease	M	3	-	-	-	-	-	-	-	3
	F	1	-	-	-	-	-	-	1	-
20. Other Heart Disease	M	5	-	-	-	-	-	-	-	5
	F	4	-	-	-	-	-	3	-	1
21. Other Circulatory Disease	M	-	-	-	-	-	-	-	-	-
	F	2	-	-	-	1	-	1	-	-
23. Pneumonia	M	2	-	-	-	-	-	-	-	2
	F	3	-	-	-	-	-	-	-	3
24. Bronchitis	M	2	-	-	-	-	-	-	-	2
	F	3	-	-	-	-	-	1	2	-
25. Other Diseases of Respiratory System	M	-	-	-	-	-	-	-	-	-
	F	1	-	-	-	-	1	-	-	-
26. Ulcer of Stomach and Duodenum	M	1	-	-	-	-	-	-	1	-

28. Nephritis and Nephrosis	M	2	-	-	-	-	-	-	-	1	1	1
	F	-	-	-	-	-	-	-	-	-	-	-
32. Other Defined and Ill-Defined Diseases	M	3	-	-	-	-	-	-	-	2	1	1
	F	3	1	-	-	-	-	-	-	-	1	1
34. All Other Accidents	M	-	-	-	-	-	-	-	-	-	-	-
	F	1	-	-	-	-	-	-	-	-	1	1
35. Suicide	M	1	-	-	-	-	-	-	-	-	-	-
	F	-	-	-	-	-	-	-	-	-	-	-
<u>TOTAL ALL CAUSES</u>	(M	34	-	-	-	-	-	-	-	5	6	23)
	F	38	1	-	-	-	-	-	-	9	7	16)

Of the deaths over 75 the ages and sex were:

	75 - 80	80 - 90	over 90
M	9	12	2
F	5	7	4

The greatest age was 93.

Raunds	Percentage of Deaths from Special Causes			
	England and Wales			
	Males & Females Combined	Males Separate	Females Separate	
Tuberculosis	0.0	0.47	0.73	0.22
Circulatory Disease	54.0	53.6	50.5	56.7
Respiratory Disease	15.0	14.0	16.3	11.6
All Cancers	14.0	18.6	19.8	17.5
Lung Cancers	1.4	4.4	7.4	1.3

ILLEGITIMATE BIRTHS

Table No. 2

RATES PER 1,000 TOTAL BIRTHS

YEAR	RAUNDS	OUNCLE & THRAPSTON	COUNTY
1935	00.0	36.0	36.0
1936	55.0	32.0	34.0
1937	46.0	34.6	36.0
1938	16.0	32.0	37.0
1939	54.0	45.6	37.0
1940	46.0	37.0	36.0
1941	17.0	79.4	44.0
1942	55.0	52.4	54.0
1943	84.0	64.3	69.0
1944	79.0	82.6	83.0
1945	20.0	129.2	109.0
1946	111.0	45.0	68.0
1947	20.0	60.0	55.0
1948	65.0	54.0	49.0
1949	15.0	62.9	46.0
1950	57.0	84.6	46.0
1951	60.0	80.0	50.0
1952	73.0	39.9	44.0
1953	00.0	40.34	41.0
1954	55.0	65.0	51.0
1955	16.0	51.3	45.0
1956	67.5	74.7	44.0
1957	61.0	60.8	41.0
1958	50.0	52.6	38.0
1959	33.3	43.0	41.0
1960	00.0	50.0	41.0
1961	30.0	41.6	51.0
1962	16.4	61.6	49.7
1963	16.4	43.6	54.1

Corresponding rates for Raunds since 1908 were:-

1908	33.0	1917	86.0	1926	36.0
1909	10.9	1918	16.0	1927	15.0
1910	00.0	1919	70.0	1928	53.0
1911	40.0	1920	37.0	1929	20.0
1912	25.0	1921	50.0	1930	00.0
1913	12.5	1922	54.0	1931	00.0
1914	51.7	1923	20.0	1932	00.0
1915	25.0	1924	50.0	1933	20.0
1916	37.5	1925	00.0	1934	20.0

In the whole of England and Wales in the year 1963 there were 59,048 illegitimate births or 69 out of every 1000 live births. For the previous 5 years 1962, 1961, 1960, 1959 and 1958 the rate was 56 per 1000 live births. The national illegitimate rate has thus gone up 23% in 1963.

A TABLE OF BIRTH RATES AND DEATH RATES FROM SPECIAL CAUSES SINCE THE
FORMATION OF THE DISTRICT ON 1st APRIL, 1935.

Table No. 3

DEATHS													
Estimated Population		Births		All ages		Under 1		Pulm. Tubercu.		Non-Pulm. Tubercu.		Cancer	
		No.	Rate per 1000 Pop.	No.	Rate per 1000 Pop.	No.	Rate per 1000 Births	No.	Rate per 1000 Pop.	No.	Rate per 1000 Pop.	No.	Rate per 1000 Pop.
1935	...	45	10.6	40	8.5	1	22.0	1	0.22	0	0.00	6	1.4
1936	...	62	14.0	65	13.37	3	48.0	4	0.90	0	0.00	12	2.7
1937	...	65	14.7	67	13.7	1	15.0	4	0.90	0	0.00	12	2.7
1938	...	63	14.2	49	10.07	1	16.0	4	0.90	0	0.00	5	1.1
1939	...	55	12.3	45	9.2	3	59.0	0	0.00	0	0.00	7	1.6
1940	...	65	14.21	55	10.3	2	30.0	1	0.22	0	0.00	2	0.4
1941	...	58	10.7	62	11.5	3	51.0	2	0.37	1	0.18	7	1.3
1942	...	73	14.1	48	9.8	0	0.0	4	0.77	0	0.00	9	1.7
1943	...	71	14.7	56	11.6	6	84.0	2	0.41	0	0.00	9	1.8
1944	...	85	20.7	61	14.3	3	36.0	1	0.23	1	0.23	12	2.8
1945	...	80	18.9	71	16.8	4	50.0	2	0.47	0	0.00	9	2.1
1946	...	89	20.4	53	12.16	5	56.0	1	0.23	0	0.00	7	1.6
1947	...	98	22.2	58	13.14	6	61.0	1	0.22	0	0.00	10	2.6
1948	...	77	17.18	65	14.5	5	65.0	1	0.22	0	0.00	8	1.8
1949	...	65	14.3	65	11.8	3	46.0	0	0.00	0	0.00	8	1.8
1950	...	53	12.8	64	11.6	3	57.0	0	0.00	0	0.00	6	1.3
1951	...	67	16.0	62	11.1	1	15.0	2	0.43	0	0.00	10	2.2
1952	...	55	13.1	75	13.5	2	36.0	1	0.22	0	0.00	10	2.2
1953	...	66	16.5	41	7.2	2	30.0	0	0.00	0	0.00	6	1.3
1954	...	73	18.13	44	7.41	2	27.3	0	0.00	0	0.00	3	0.6
1955	...	62	15.46	81	13.6	2	27.3	1	0.21	0	0.00	17	3.6
1956	...	74	18.5	56	10.05	1	13.5	1	0.21	0	0.00	11	2.35
1957	...	82	20.41	52	9.48	2	24.4	0	0.00	0	0.00	10	2.14
1958	...	60	14.19	52	9.5	0	0.0	1	0.21	0	0.00	8	1.72
1959	...	60	14.87	62	11.12	1	16.6	0	0.00	0	0.00	9	1.92
1960	...	56	13.92	56	10.3	1	17.86	0	0.00	0	0.00	10	2.14
1961	...	67	17.0	58	10.53	3	46.0	0	0.00	0	0.00	12	2.62
1962	...	61	15.3	64	11.8	1	16.1	0	0.00	0	0.00	11	2.38
1963	...	61	15.0	72	13.1	1	16.1	0	0.00	0	0.00	10	2.46
TOTAL FOR 29 YEARS :-		1948		1699		68		34		2		256	
Raunds Urban District													

Birth Rate is Standard Rate from 1950 onwards. Death Rate is Standard Rate except for 1942-1948, both inclusive.

A TABLE OF BIRTH RATES AND DEATH RATES FROM SPECIAL CAUSES 1899 - 1934

Table No. 4

		Births		DEATHS											
				All ages		Under 1		Pulm. Tubercu.		Non-Pulm. Tubercu.		Cancer			
								No.	Rate per 1000 Pop.	No.	Rate per 1000 Pop.	No.	Rate per 1000 Pop.	No.	Rate per 1000 Pop.
Estimated Population		No.	Crude Rate per 1000 Pop.	No.	Crude Rate per 1000 Pop.	No.	Rate per 1000 Births	No.	Rate per 1000 Pop.	No.	Rate per 1000 Pop.	No.	Rate per 1000 Pop.	No.	Rate per 1000 Pop.
1899	...	109	28.6	38	9.9	11	109								
1900	...	123	32.2	42	11.0	10	81								
1901	...	118	30.2	44	11.2	7	59								
1902	...	115	28.5	48	11.9	12	104								
1903	...	109	25.8	45	10.6	14	128								
1904	...	109	24.9	47	10.7	13	119								
1905	...	101	23.0	36	8.2	10	99								
1906	...	103	23.5	42	9.5	8	77.6								
1907	...	99	23.7	47	10.9	9	91								
1908	...	120	27.8	52	12.0	13	108	8	1.85	2	0.46	2	0.46	2	0.46
1909	...	91	21.6	40	9.2	4	44	6	1.39	2	0.46	5	0.46	5	0.46
1910	...	78	18.0	47	10.9	8	102	7	1.61	6	1.38	4	1.38	6	1.38
1911	...	76	19.6	50	12.9	5	65	4	1.03	4	1.03	2	1.03	2	1.03
1912	...	80	20.6	35	9.0	4	50	3	0.77	6	0.50	2	0.50	5	0.50
1913	...	81	20.9	51	13.1	7	86	7	1.80	6	1.54	5	1.54	5	1.54
1914	...	58	14.9	47	12.1	5	86	5	1.30	4	1.05	5	1.05	5	1.05
1915	...	80	19.7	51	12.5	9	112	2	0.50	4	1.00	6	1.00	6	1.00
1916	...	80	19.7	48	11.7	7	87	5	1.22	2	0.50	5	0.50	5	0.50
1917	Births Deaths	62	15.5	41	11.1	2	32	6	1.66	3	0.83	2	0.83	2	0.83
1918	Births Deaths	62	14.6	54	14.3	3	48	5	1.325	5	1.326	4	1.326	4	1.326
1919	Births Deaths	57	13.3	40	9.9	2	35	2	0.49	0	0.0	7	0.0	7	0.0
1920	...	80	19.0	45	10.7	5	62	4	0.936	1	0.234	5	0.234	5	0.234
1921	...	62	16.2	39	10.2	3	48	2	0.52	1	0.26	2	0.26	2	0.26
1922	...	55	14.3	34	8.8	3	54	6	1.5	0	0.0	1	0.0	1	0.0
1923	...	53	13.9	46	12.0	7	134	5	1.3	2	0.52	2	0.52	2	0.52
1924	...	60	15.6	41	10.7	5	83	3	0.78	2	0.52	6	0.52	6	0.52
1925	...	53	13.8	32	8.3	1	18	5	1.3	0	0.0	4	0.0	4	0.0

1926	...	...	3753	55	14.6	50	13.3	2	36	1	0.26	0	0.0	6
1927	...	...	3690	41	11.1	37	10.0	3	73	1	0.26	0	0.0	0
1928	...	...	3715	56	15.0	32	8.6	0	00	2	0.54	2	0.54	4
1929	...	...	3753	46	12.2	48	12.7	4	87	5	1.35	1	0.27	3
1930	...	...	3753	41	10.9	34	9.0	0	00	3	0.798	1	0.266	2
1931	...	...	3687	37	10.0	40	10.8	1	27	2	0.54	1	0.27	8
1932	...	...	3626	37	10.2	48	13.2	5	135	1	0.27	0	0.0	7
1933	...	...	3561	46	12.9	48	13.4	2	44	5	1.4	0	0.0	6
1934	...	...	3617	52	14.3	45	11.3	7	135	1	0.27	1	0.27	8
TOTAL FOR 36 YEARS :- 1899 - 1934 Rounds Urban District				2685		1568		212		107 in 27 yrs.		50 in 27 yrs.		118 in 27 yrs.

The table below tells its own story, but it would be a delusion to conclude that cancer is on the increase. The apparent increase is due to more exact diagnosis. The Infantile death rate and the Tuberculosis death rate are related to social and hygenic conditions.

	Mortality Under 1 Year		Mortality Pulm.Tubercu.		Mortality Cancer	
	Total No.	Rate	Total No.	Rate	Total No.	Rate
		per 1000 Births		per 1000 Pop.		per 1000 Pop.
Raunds Urban District						
1899-1908	107	97.0				
1909-1918	54	72.0	53	1.3	43	
1919-1928	31	54.0	34	0.9	37	
1936-1945	26	38.9	24	0.517	84	
1952-1961	16	24.43	4	0.085	96	
Administrative County						
1936-1945	1560	42.15	1028	0.44		
1952-1961	993	21.63	253	0.09		

Natural Increase of population is Birth numbers - Death numbers which, in 1963, was 72 - 61 = 11. It should be understood that Standard Rates are for comparison with the rest of England and Wales, whereas Crude Rates are the actual rate. From the foregoing Table, Standard Births : Deaths are 16 : 13.1, whereas the Crude Rates are 13.2 : 15.6, which means a reduction in population of 4,610 of 11. The natural increase in the whole of England and Wales for 1962 was 283,111; in 1961 an increase of 252,900 and during the years 1956- 1960 an average increase of 215,311. The total population of England and Wales at 30th June, 1962 was estimated to be 46,669,000 made up of 22,651,000 males and 24,018,000 females. In 1963 there was an estimated population of 47,023,000 made up of 22,832,000 males and 24,191,000 females.

It may be of general interest to know that the total population of 47,023,000 at 30th June, 1963 was according to age groups and sex, made up of:

	All Ages	0-	5-	15-	25-	35-	45-	55-	65-	75
Males	22,832	2,001	3,474	3,348	3,007	3,203	3,040	2,612	1,463	684
Females	24,191	1,898	3,296	3,284	2,886	3,201	3,163	2,960	2,163	1,340

These figures are expressed in number of thousands. Evidently the males predominate until between the ages of 25 to 35, after which the females predominate. Cver 70 females : males are as 3 : 2; at 80 the rate is 2 : 1.

The groups between the working ages of 15 - 60 years represent 60% of the total; those over 60 represent 17.4% of the total. Males over 65 (pensionable age) represent 9.4% of all males and females over 60 (pensionable age) represent 20.2% of all females. Combined males and females of pensionable age represent 15% of the total population.

The comparative table below shows the decrease or increase in decades of life of males and females; the number at 5 - 15 years being numcrically 100, and this is related to the number of each sex at 5 - 15 years.

			Males	Females
0 - 5 years	...	...	115	115
5 - 15 "	...	...	100	100
15 - 25 "	...	...	96.0	99.0
25 - 35 "	...	...	86.0	87.5
35 - 45 "	...	...	92.0	97.0
45 - 55 "	...	...	87.0	96.0
55 - 65 "	...	...	75.0	90.0
65 - 75 "	...	...	42.0	65.6
75 & over	...	...	19.6	40.0

